

# Refined Liouville-Type Results for the Three-Dimensional Stationary MHD Equations

## Abstract

We establish refined Liouville-type theorems for smooth solutions to the three-dimensional stationary magnetohydrodynamic (MHD) equations. By introducing a general growth function  $g(\rho)$  that only satisfies mild conditions (monotonicity,  $\rho^{-1/3}g(\rho) \rightarrow 0$ , and  $\int^\infty \frac{d\rho}{\rho g(\rho)} = \infty$ ), we prove that any solution where the velocity  $\mathbf{u}$  and magnetic field  $\mathbf{b}$  grow at most like  $\rho^{\frac{2}{p}-\frac{1}{3}}g(\rho)^{\frac{3}{p}-1}$  (for some  $3/2 < p < 3$ ) must be identically zero. The proof combines local energy estimates for the coupled MHD system, the Bogovskiĭ operator to handle the pressure, an iterative scheme, and a contradiction argument based on the divergence of the integral of  $1/(\rho g(\rho))$ . The result extends recent sharp Liouville theorems for the Navier–Stokes equations to the MHD case and allows for logarithmic or even weaker subcritical growth.

**Keywords:** MHD equations, Liouville-type theorem, energy estimates

**MSC(2020):** 35Q35, 35B53, 76W05

## 1 Introduction

We consider smooth solutions to the three-dimensional stationary incompressible MHD equations

$$\begin{cases} -\Delta \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} - (\mathbf{b} \cdot \nabla) \mathbf{b} + \nabla \pi = 0, \\ -\Delta \mathbf{b} + (\mathbf{u} \cdot \nabla) \mathbf{b} - (\mathbf{b} \cdot \nabla) \mathbf{u} = 0, \\ \nabla \cdot \mathbf{u} = \nabla \cdot \mathbf{b} = 0, \end{cases} \quad (1)$$

where  $\mathbf{u} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  denotes the velocity field,  $\mathbf{b} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  the magnetic field, and  $\pi : \mathbb{R}^3 \rightarrow \mathbb{R}$  the pressure. The system describes the steady motion of an electrically conducting fluid in the absence of external forces. A classical question is to determine under which decay or growth conditions the only smooth solution is the trivial one,  $\mathbf{u} \equiv 0$ ,  $\mathbf{b} \equiv 0$ ; results of this type are known as Liouville-type theorems.

For the stationary Navier-Stokes equations ( $\mathbf{b} \equiv 0$ ), Liouville theorems have a long history. Galdi [1] proved that if  $\mathbf{u} \in L^{9/2}(\mathbb{R}^3)$  then  $\mathbf{u} \equiv 0$ . Later, Seregin [2] treated the case when the velocity field can be written as  $\mathbf{u} = \nabla \cdot V$  with a skew-symmetric  $V \in \text{BMO}(\mathbb{R}^3)$ . Chae and Wolf [5, 6] extended this framework to growth conditions on

the potential  $V$  over large balls. In a different direction, Seregin and Wang [3] employed Lorentz spaces to obtain decay criteria, showing that

$$\int_{\mathbb{R}^3} |\nabla \mathbf{u}|^2 \leq C \liminf_{\rho \rightarrow \infty} \frac{\|\mathbf{u}\|_{L^{p,l}(A(\rho, 2\rho))}}{\rho^{\frac{9}{p}-2}}$$

for  $p > 3$ , which implies triviality under suitable smallness. Tsai [4] proved that  $\mathbf{u} \equiv 0$  if

$$\liminf_{\rho \rightarrow \infty} \frac{\|\mathbf{u}\|_{L^p(A(\rho, 2\rho))}}{\rho^{\frac{2}{p}-\frac{1}{3}}} = 0 \quad \text{for some } \frac{12}{5} \leq p \leq 3.$$

Cho, Neustupa and Yang [7] sharpened this condition by showing that it suffices that the liminf is finite (instead of zero) and that the range can be extended to  $3/2 < p < 3$ . Their result also covered the MHD and Hall-MHD equations, establishing that the growth rate  $\rho^{\frac{2}{p}-\frac{1}{3}}$  is essentially the critical threshold for Liouville theorems in this family.

More recently, further refinements have been obtained by introducing additional logarithmic or slowly varying factors. Bang and Yang [10] studied the stationary Navier–Stokes equations and obtained Saint-Venant type estimates with logarithmic corrections. These results indicate that the sharp condition can be weakened by allowing an arbitrarily slowly diverging multiplicative factor  $g(\rho)$ . The purpose of the present paper is to systematically generalize the Liouville theorem for the MHD equations by introducing a general growth function  $g(\rho)$  satisfying mild conditions (see Assumption 1.3 below). We show that if the  $L^p$  norms of  $\mathbf{u}$  and  $\mathbf{b}$  over large annuli are bounded by  $\rho^{\frac{2}{p}-\frac{1}{3}}g(\rho)^{\frac{3}{p}-1}$  for some  $3/2 < p < 3$ , then  $\mathbf{u} = \mathbf{b} \equiv 0$ . The function  $g$  is only required to be non-decreasing, to satisfy  $\rho^{-1/3}g(\rho) \rightarrow 0$ , and to make the integral  $\int^\infty \frac{d\rho}{\rho g(\rho)}$  diverge. Typical examples include iterated logarithms such as  $g(\rho) = \log(e + \rho)$ ,  $\log(e + \log(e + \rho))$ , etc., as well as certain power-like functions constructed by induction.

Compared to the pure Navier-Stokes case, the MHD system (1) presents additional difficulties due to the coupling terms between  $\mathbf{u}$  and  $\mathbf{b}$ . We treat them by careful integration by parts, which transforms the Lorentz force and the magnetic transport terms into boundary integrals that can be estimated exactly as the convective terms. This allows us to transfer the energy estimates developed for the Navier-Stokes equations to the MHD setting. Our main results read as follows.

**Theorem 1.1.** *Let  $g$  satisfy Assumption 1.3 below. Suppose  $(\mathbf{u}, \mathbf{b}, \pi)$  is a smooth solution to (1) such that*

$$\nabla \mathbf{u}, \nabla \mathbf{b} \in L^2(\mathbb{R}^3),$$

*and that there exist  $0 < \theta < 1$  and  $1 \leq p < 3$  satisfying*

$$\limsup_{\rho \rightarrow \infty} \frac{\|\mathbf{u}\|_{L^p(A(\theta\rho, \theta^{-1}\rho))} + \|\mathbf{b}\|_{L^p(A(\theta\rho, \theta^{-1}\rho))}}{\rho^{\frac{2}{p}-\frac{1}{3}}g(\rho)^{\frac{3}{p}-1}} < \infty. \quad (2)$$

*Then  $\mathbf{u} = \mathbf{b} \equiv 0$ .*

**Theorem 1.2.** *Let  $g$  satisfy Assumption 1.3. Suppose  $(\mathbf{u}, \mathbf{b}, \pi)$  is a smooth solution to (1) such that there exist  $0 < \theta < 1$  and  $\frac{3}{2} < p < 3$  satisfying (2). Then  $\mathbf{u} = \mathbf{b} \equiv 0$ .*

In Theorem 1.2 the a priori condition  $\nabla \mathbf{u}, \nabla \mathbf{b} \in L^2$  is not required; it follows from the growth restriction and the equations. The range  $3/2 < p < 3$  is the same as in the Navier–Stokes case and is believed to be optimal within the energy framework.

The function  $g$  must satisfy:

*Assumption 1.3.* A function  $g : [1, \infty) \rightarrow [1, \infty)$  satisfies

$$g \text{ is non-decreasing on } [1, \infty), \lim_{\rho \rightarrow \infty} \rho^{-1/3} g(\rho) = 0, \text{ for all } a \geq 1, \int_a^\infty \frac{1}{g(\rho)} \frac{d\rho}{\rho} = \infty.$$

Examples of admissible  $g$  are iterated logarithms or functions constructed by gluing powers (see e.g. [7] for a detailed explanation).

## 2 Preliminaries

We denote  $B(\rho) = \{x \in \mathbb{R}^3 : |x| < \rho\}$  and  $A(\rho_1, \rho_2) = \{x : \rho_1 \leq |x| < \rho_2\}$  for  $0 < \rho_1 < \rho_2 < \infty$ . Generic constants are written as  $C$  and may depend on  $\theta, p$ , etc. We write  $a \lesssim b$  if  $a \leq Cb$ , and  $a \lesssim_q b$  if the constant depends on  $q$ . The average of  $f$  over a measurable set  $\Omega$  is  $(f)_\Omega = \frac{1}{|\Omega|} \int_\Omega f$ .

For  $1 < q < \infty$ ,  $L_0^q(\Omega) = \{f \in L^q(\Omega) : (f)_\Omega = 0\}$ , and  $W_0^{1,q}(\Omega)$  is the closure of  $C_c^\infty(\Omega)$  in the Sobolev space  $W^{1,q}(\Omega)$ . A fundamental tool is the Bogovskiĭ operator providing a right inverse of the divergence on annular domains.

**Lemma 2.1** ([4, Lemma 3]). *Let  $0 < \theta < 1$ ,  $R > 1$ , and  $1 < q < \infty$ . There exists a linear map*

$$T : L_0^q(A(\theta R, R)) \rightarrow W_0^{1,q}(A(\theta R, R))$$

*such that for any  $f \in L_0^q(A(\theta R, R))$ ,*

$$\nabla \cdot Tf = f, \quad \|\nabla Tf\|_{L^q(A(\theta R, R))} \leq C_{\theta,q} \|f\|_{L^q(A(\theta R, R))}.$$

*Moreover,  $C_{\theta,q} \rightarrow \infty$  as  $\theta \rightarrow 1$ .*

Fix  $0 < \theta < 1$  and define the piecewise linear cut-off  $\eta_\theta : [0, \infty) \rightarrow [0, 1]$  by

$$\eta_\theta(t) = \begin{cases} 1, & 0 \leq t \leq \theta, \\ \frac{t - \theta}{1 - \theta}, & \theta < t < 1, \\ 0, & t \geq 1. \end{cases}$$

The associated energy functional for the MHD system is

$$E_\theta(\rho) = \int_{\mathbb{R}^3} (|\nabla \mathbf{u}(x)|^2 + |\nabla \mathbf{b}(x)|^2) \eta_\theta\left(\frac{|x|}{\rho}\right) dx, \quad \rho > 0.$$

**Lemma 2.2.**  *$E_\theta \in C^1((0, \infty))$  and for all  $\rho > 0$*

$$E'_\theta(\rho) = \frac{1}{\rho^2(1 - \theta)} \int_{A(\theta\rho, \rho)} (|\nabla \mathbf{u}|^2 + |\nabla \mathbf{b}|^2) |x| dx.$$

*In particular,*

$$E'_\theta(\rho) \geq \frac{\theta}{\rho(1 - \theta)} (\|\nabla \mathbf{u}\|_{L^2(A(\theta\rho, \rho))}^2 + \|\nabla \mathbf{b}\|_{L^2(A(\theta\rho, \rho))}^2).$$

- 3. Proof.** The proof follows the standard differentiation of piecewise linear cut-offs. Because  $\eta_\theta$  is constant outside  $(\theta, 1)$ , the difference quotient  $\frac{1}{\rho h} [E_\theta(\rho(1 + h)) - E_\theta(\rho)]$  reduces to integrals over the transition regions. The linearity of  $\eta_\theta$  yields the exact expression for  $E'_\theta(\rho)$ ; the lower bound follows because  $|x| \geq \theta\rho$  on  $A(\theta\rho, \rho)$ . See, e.g., [4] for the scalar case.  $\square$

### 3 Energy estimates for the MHD system

Throughout this section we fix  $0 < \theta < 1$  and let  $\rho \geq 1$ . Define

$$K_\theta(\rho) = \rho^{-1} (\|\mathbf{u}\|_{L^3(A(\theta\rho, \theta^{-1}\rho))}^3 + \|\mathbf{b}\|_{L^3(A(\theta\rho, \theta^{-1}\rho))}^3).$$

#### 3.1 Basic local energy inequality

**Lemma 3.1.** *[label=()]*

1. If  $\liminf_{\rho \rightarrow \infty} K_\theta(\rho) = 0$ , then  $\mathbf{u} = \mathbf{b} \equiv 0$ .
2. If  $\liminf_{\rho \rightarrow \infty} K_\theta(\rho) < \infty$ , then  $\nabla \mathbf{u}, \nabla \mathbf{b} \in L^2(\mathbb{R}^3)$  and  $\mathbf{u}, \mathbf{b} \in L^6(\mathbb{R}^3)$ .

*Proof.* Let  $\rho \leq r < R \leq \theta^{-1}\rho$  and choose a radially decreasing  $\phi \in C_c^2(B(R))$  with  $\phi \equiv 1$  on  $B(r)$  and

$$|\nabla \phi| \leq \frac{C}{R-r}, \quad |\nabla^2 \phi| \leq \frac{C}{(R-r)^2},$$

where  $C$  is an absolute constant. Set  $S = A(\theta R, R) \subset A(\theta\rho, \theta^{-1}\rho)$ .

Since  $\nabla \cdot \mathbf{u} = 0$ ,  $\mathbf{u} \cdot \nabla \phi^2$  has zero mean over  $S$ . By Lemma 2.1 we obtain  $\mathbf{v} = T(\mathbf{u} \cdot \nabla \phi^2) \in W_0^{1,q}(S)$  satisfying  $\nabla \cdot \mathbf{v} = \mathbf{u} \cdot \nabla \phi^2$  and, for any  $1 < q < \infty$ ,

$$\|\nabla \mathbf{v}\|_{L^q(S)} \lesssim_q \frac{1}{R-r} \|\mathbf{u}\|_{L^q(S)}. \quad (3)$$

Extend  $\mathbf{v}$  by zero outside  $S$ .

Testing the first equation of (1) with  $\mathbf{u}\phi^2 - \mathbf{v}$  and integrating over  $\mathbb{R}^3$  yields

$$\int -\Delta \mathbf{u} \cdot (\mathbf{u}\phi^2 - \mathbf{v}) + \int (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot (\mathbf{u}\phi^2 - \mathbf{v}) - \int (\mathbf{b} \cdot \nabla) \mathbf{b} \cdot (\mathbf{u}\phi^2 - \mathbf{v}) + \int \nabla \pi \cdot (\mathbf{u}\phi^2 - \mathbf{v}) = 0.$$

Since  $\nabla \cdot \mathbf{u} = 0$ , the pressure term vanishes after integration by parts:

$$\int \nabla \pi \cdot (\mathbf{u}\phi^2 - \mathbf{v}) = - \int \pi \nabla \cdot (\mathbf{u}\phi^2 - \mathbf{v}) = 0.$$

For the other terms we integrate by parts repeatedly. The Laplacian term gives

$$\int -\Delta \mathbf{u} \cdot (\mathbf{u}\phi^2 - \mathbf{v}) = \int |\nabla \mathbf{u}|^2 \phi^2 - \frac{1}{2} \int |\mathbf{u}|^2 \Delta \phi^2 + \int \nabla \mathbf{u} : (\nabla \mathbf{v})^T.$$

The convection term rewrites as

$$\begin{aligned} \int (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot (\mathbf{u}\phi^2 - \mathbf{v}) &= \frac{1}{2} \int \mathbf{u} \cdot \nabla |\mathbf{u}|^2 \phi^2 - \int (\mathbf{u} \cdot \nabla) \mathbf{u} \cdot \mathbf{v} \\ &= -\frac{1}{2} \int |\mathbf{u}|^2 \mathbf{u} \cdot \nabla \phi^2 - \int (\mathbf{u} \otimes \mathbf{u}) : \nabla \mathbf{v}, \end{aligned}$$

where we integrated by parts in the first part and used  $\nabla \cdot \mathbf{u} = 0$ . The Lorentz term is kept as

$$\int (\mathbf{b} \cdot \nabla) \mathbf{b} \cdot (\mathbf{u}\phi^2 - \mathbf{v}).$$

Collecting, the momentum equation gives

$$\begin{aligned} \int |\nabla \mathbf{u}|^2 \phi^2 &= \frac{1}{2} \int |\mathbf{u}|^2 \Delta \phi^2 + \int \nabla \mathbf{u} : (\nabla \mathbf{v})^T \\ &+ \frac{1}{2} \int |\mathbf{u}|^2 \mathbf{u} \cdot \nabla \phi^2 - \int (\mathbf{u} \otimes \mathbf{u}) : (\nabla \mathbf{v})^T \\ &+ \int (\mathbf{b} \cdot \nabla) \mathbf{b} \cdot (\mathbf{u} \phi^2 - \mathbf{v}). \end{aligned} \quad (4)$$

Testing the second equation of (1) with  $\mathbf{b} \phi^2$  yields

$$\int -\Delta \mathbf{b} \cdot \mathbf{b} \phi^2 + \int (\mathbf{u} \cdot \nabla) \mathbf{b} \cdot \mathbf{b} \phi^2 - \int (\mathbf{b} \cdot \nabla) \mathbf{u} \cdot \mathbf{b} \phi^2 = 0.$$

The Laplacian term gives  $\int |\nabla \mathbf{b}|^2 \phi^2 - \frac{1}{2} \int |\mathbf{b}|^2 \Delta \phi^2$ . For the advection term we integrate by parts using  $\nabla \cdot \mathbf{u} = 0$ :

$$\int (\mathbf{u} \cdot \nabla) \mathbf{b} \cdot \mathbf{b} \phi^2 = -\frac{1}{2} \int |\mathbf{b}|^2 \nabla \cdot (\mathbf{u} \phi^2) = -\frac{1}{2} \int |\mathbf{b}|^2 \mathbf{u} \cdot \nabla \phi^2.$$

The remaining term is kept. Thus

$$\int |\nabla \mathbf{b}|^2 \phi^2 = \frac{1}{2} \int |\mathbf{b}|^2 \Delta \phi^2 + \frac{1}{2} \int |\mathbf{b}|^2 \mathbf{u} \cdot \nabla \phi^2 + \int (\mathbf{b} \cdot \nabla) \mathbf{u} \cdot \mathbf{b} \phi^2. \quad (5)$$

Adding (4) and (5) we obtain the total energy identity

$$\begin{aligned} \int (|\nabla \mathbf{u}|^2 + |\nabla \mathbf{b}|^2) \phi^2 &= \frac{1}{2} \int (|\mathbf{u}|^2 + |\mathbf{b}|^2) \Delta \phi^2 + \int \nabla \mathbf{u} : (\nabla \mathbf{v})^T \\ &+ \frac{1}{2} \int (|\mathbf{u}|^2 - |\mathbf{b}|^2) \mathbf{u} \cdot \nabla \phi^2 - \int (\mathbf{u} \otimes \mathbf{u}) : (\nabla \mathbf{v})^T \\ &+ \underbrace{\int (\mathbf{b} \cdot \nabla) \mathbf{b} \cdot \mathbf{u} \phi^2 + \int (\mathbf{b} \cdot \nabla) \mathbf{u} \cdot \mathbf{b} \phi^2 - \int (\mathbf{b} \cdot \nabla) \mathbf{b} \cdot \mathbf{v}}_{=: I_{\text{coup}}}. \end{aligned}$$

For the first two coupling terms we integrate by parts:

$$\int (\mathbf{b} \cdot \nabla) \mathbf{b} \cdot \mathbf{u} \phi^2 = - \int (\mathbf{b} \otimes \mathbf{b}) : \nabla \mathbf{u} \phi^2 - \int (\mathbf{b} \otimes \mathbf{b}) : (\mathbf{u} \otimes \nabla \phi^2),$$

while

$$\int (\mathbf{b} \cdot \nabla) \mathbf{u} \cdot \mathbf{b} \phi^2 = \int (\mathbf{b} \otimes \mathbf{b}) : \nabla \mathbf{u} \phi^2.$$

Summing them cancels the volume term involving  $\nabla \mathbf{u}$ , leaving

$$\int (\mathbf{b} \cdot \nabla) \mathbf{b} \cdot \mathbf{u} \phi^2 + \int (\mathbf{b} \cdot \nabla) \mathbf{u} \cdot \mathbf{b} \phi^2 = - \int (\mathbf{b} \otimes \mathbf{b}) : (\mathbf{u} \otimes \nabla \phi^2).$$

The third term is handled similarly:

$$\int (\mathbf{b} \cdot \nabla) \mathbf{b} \cdot \mathbf{v} = - \int (\mathbf{b} \otimes \mathbf{b}) : \nabla \mathbf{v}.$$

Therefore

$$\begin{aligned} \int (|\nabla \mathbf{u}|^2 + |\nabla \mathbf{b}|^2) \phi^2 &= \frac{1}{2} \int (|\mathbf{u}|^2 + |\mathbf{b}|^2) \Delta \phi^2 + \int \nabla \mathbf{u} : (\nabla \mathbf{v})^T \\ &\quad + \frac{1}{2} \int (|\mathbf{u}|^2 - |\mathbf{b}|^2) \mathbf{u} \cdot \nabla \phi^2 - \int (\mathbf{u} \otimes \mathbf{u}) : (\nabla \mathbf{v})^T \\ &\quad - \int (\mathbf{b} \otimes \mathbf{b}) : (\mathbf{u} \otimes \nabla \phi^2) + \int (\mathbf{b} \otimes \mathbf{b}) : \nabla \mathbf{v}. \end{aligned} \quad (6)$$

Denote the six terms on the right by  $J_1 - J_6$ .

Using Hölder's inequality, the bounds on  $\phi$  and (3), we get

$$\begin{aligned} J_1 &\lesssim (R-r)^{-2} (\|\mathbf{u}\|_{L^2(S)}^2 + \|\mathbf{b}\|_{L^2(S)}^2), \\ J_2 &\leq \|\nabla \mathbf{u}\|_{L^2(S)} \|\nabla \mathbf{v}\|_{L^2(S)} \lesssim \frac{1}{R-r} \|\nabla \mathbf{u}\|_{L^2(S)} \|\mathbf{u}\|_{L^2(S)}, \\ J_3 &\lesssim \frac{1}{R-r} (\|\mathbf{u}\|_{L^3(S)}^3 + \|\mathbf{b}\|_{L^3(S)}^2 \|\mathbf{u}\|_{L^3(S)}), \\ J_4 &\leq \|\mathbf{u}\|_{L^3(S)}^2 \|\nabla \mathbf{v}\|_{L^3(S)} \lesssim \frac{1}{R-r} \|\mathbf{u}\|_{L^3(S)}^3, \\ J_5 &\lesssim \frac{1}{R-r} \|\mathbf{b}\|_{L^3(S)}^2 \|\mathbf{u}\|_{L^3(S)}, \\ J_6 &\leq \|\mathbf{b}\|_{L^3(S)}^2 \|\nabla \mathbf{v}\|_{L^3(S)} \lesssim \frac{1}{R-r} \|\mathbf{b}\|_{L^3(S)}^2 \|\mathbf{u}\|_{L^3(S)}. \end{aligned}$$

For  $J_2$  we apply Young's inequality: for any  $\epsilon > 0$ ,

$$J_2 \leq \epsilon \|\nabla \mathbf{u}\|_{L^2(S)}^2 + \frac{C_\epsilon}{(R-r)^2} \|\mathbf{u}\|_{L^2(S)}^2.$$

By Hölder and  $|S| \lesssim R^3 \leq (\theta^{-1}\rho)^3$ , we have  $\|\mathbf{u}\|_{L^2(S)}^2 \lesssim \rho \|\mathbf{u}\|_{L^3(S)}^2$ . Then Young's inequality again gives

$$\frac{\rho}{(R-r)^2} \|\mathbf{u}\|_{L^3(S)}^2 \leq \frac{\|\mathbf{u}\|_{L^3(S)}^3}{R-r} + \frac{\rho^3}{(R-r)^4}.$$

All other  $J_k$  ( $k = 3, \dots, 6$ ) are directly bounded by  $\frac{1}{R-r} (\|\mathbf{u}\|_{L^3(S)}^3 + \|\mathbf{b}\|_{L^3(S)}^3)$  up to constants.

Collecting these estimates and absorbing the  $\epsilon \|\nabla \mathbf{u}\|_{L^2(S)}^2$  into the left side, we obtain for any  $\epsilon > 0$  a constant  $C_{\epsilon, \theta}$  such that

$$\begin{aligned} &\|\mathbf{u}\|_{L^6(B(r))}^2 + \|\mathbf{b}\|_{L^6(B(r))}^2 + \|\nabla \mathbf{u}\|_{L^2(B(r))}^2 + \|\nabla \mathbf{b}\|_{L^2(B(r))}^2 \\ &\leq \epsilon (\|\nabla \mathbf{u}\|_{L^2(B(R))}^2 + \|\nabla \mathbf{b}\|_{L^2(B(R))}^2) \\ &\quad + C_{\epsilon, \theta} \left[ \frac{\|\mathbf{u}\|_{L^3(S)}^3 + \|\mathbf{b}\|_{L^3(S)}^3}{R-r} + \frac{\rho^3}{(R-r)^4} \right]. \end{aligned} \quad (7)$$

(The  $L^6$  terms appear after applying the Sobolev inequality to  $\|\mathbf{u}\phi\|_{L^6}$  and using  $|\nabla \phi| \lesssim (R-r)^{-1}$ .)

Define

$$Q(t) = \|\mathbf{u}\|_{L^6(B(t))}^2 + \|\mathbf{b}\|_{L^6(B(t))}^2 + \|\nabla \mathbf{u}\|_{L^2(B(t))}^2 + \|\nabla \mathbf{b}\|_{L^2(B(t))}^2, \quad t > 0.$$

Fix  $\epsilon = 1/4$ . Set  $r_0 = \rho$  and for  $k \geq 1$ ,

$$r_k = r_{k-1} + \frac{R - \rho}{2^k}, \quad R = \theta^{-1}\rho.$$

Applying (7) with  $r = r_{k-1}$ ,  $R = r_k$  (so that  $S \subset A(\theta\rho, \theta^{-1}\rho)$ ) gives

$$Q(r_{k-1}) \leq \frac{1}{4}Q(r_k) + C \left[ \frac{K}{r_k - r_{k-1}} + \frac{\rho^3}{(r_k - r_{k-1})^4} \right],$$

where  $K = \|\mathbf{u}\|_{L^3(A(\theta\rho, \theta^{-1}\rho))}^3 + \|\mathbf{b}\|_{L^3(A(\theta\rho, \theta^{-1}\rho))}^3$ . Iterating down to  $k = 0$ ,

$$Q(\rho) \leq \sum_{k=1}^{\infty} \left(\frac{1}{4}\right)^{k-1} C \left[ \frac{K}{r_k - r_{k-1}} + \frac{\rho^3}{(r_k - r_{k-1})^4} \right].$$

Since  $r_k - r_{k-1} = (R - \rho)/2^k$ , we have

$$Q(\rho) \lesssim \frac{K}{R - \rho} + \frac{\rho^3}{(R - \rho)^4}.$$

Substituting  $R = \theta^{-1}\rho$  yields

$$Q(\rho) \lesssim K_\theta(\rho) + \rho^{-1}. \tag{8}$$

If  $\liminf_{\rho \rightarrow \infty} K_\theta(\rho) = 0$ , choose a subsequence  $\rho_j \rightarrow \infty$  with  $K_\theta(\rho_j) \rightarrow 0$ ; then  $Q(\rho_j) \rightarrow 0$ , forcing  $\mathbf{u} = \mathbf{b} = 0$  globally. If  $\liminf_{\rho \rightarrow \infty} K_\theta(\rho) < \infty$ , then  $Q(\rho)$  is bounded uniformly in  $\rho$ , which implies  $\nabla \mathbf{u}, \nabla \mathbf{b} \in L^2(\mathbb{R}^3)$  and  $\mathbf{u}, \mathbf{b} \in L^6(\mathbb{R}^3)$ .  $\square$

### 3.2 Improved estimate for $3/2 < p < 3$

**Lemma 3.2.** *Let  $0 < \theta < 1$  and  $\frac{3}{2} < p < 3$ . Then for all  $\rho > 1$ ,*

$$\begin{aligned} & \|\mathbf{u}\|_{L^6(B(\rho))}^2 + \|\mathbf{b}\|_{L^6(B(\rho))}^2 + \|\nabla \mathbf{u}\|_{L^2(B(\rho))}^2 + \|\nabla \mathbf{b}\|_{L^2(B(\rho))}^2 \\ & \leq C_\theta \rho^{-\frac{6-p}{2p-3}} (\|\mathbf{u}\|_{L^p(A)} + \|\mathbf{b}\|_{L^p(A)})^{\frac{3p}{2p-3}} + C_\theta \rho^{-1}, \end{aligned}$$

where  $A = A(\theta\rho, \theta^{-1}\rho)$ .

*Proof.* We start from (7). The  $L^3$  norms can be interpolated between  $L^p$  and  $L^6$ :

$$\|f\|_{L^3} \leq \|f\|_{L^p}^{\frac{p}{6-p}} \|f\|_{L^6}^{\frac{6-2p}{6-p}}, \quad \frac{3}{2} < p < 3.$$

Consequently, for any  $\eta > 0$ ,

$$\|f\|_{L^3}^3 \leq \eta \|f\|_{L^6}^2 + C_\eta \|f\|_{L^p}^{\frac{3p}{2p-3}}.$$

Apply this to  $\mathbf{u}$  and  $\mathbf{b}$  inside the term  $\frac{\|\mathbf{u}\|_{L^3(S)}^3 + \|\mathbf{b}\|_{L^3(S)}^3}{R-r}$ . We obtain

$$\frac{\|\mathbf{u}\|_{L^3(S)}^3 + \|\mathbf{b}\|_{L^3(S)}^3}{R-r} \leq \frac{2\eta}{R-r} (\|\mathbf{u}\|_{L^6(S)}^2 + \|\mathbf{b}\|_{L^6(S)}^2) + \frac{C_\eta}{R-r} (\|\mathbf{u}\|_{L^p(S)}^{\frac{3p}{2p-3}} + \|\mathbf{b}\|_{L^p(S)}^{\frac{3p}{2p-3}}).$$

Insert this into (7) and choose  $\eta$  sufficiently small so that the  $L^6$  terms can be absorbed after iteration (exactly as in Lemma 3.1). The iteration then yields

$$Q(\rho) \lesssim \frac{\|\mathbf{u}\|_{L^p(A)}^{\frac{3p}{2p-3}} + \|\mathbf{b}\|_{L^p(A)}^{\frac{3p}{2p-3}}}{(R - \rho)^{\frac{6-p}{2p-3}}} + \frac{\rho^3}{(R - \rho)^4}.$$

Setting  $R = \theta^{-1}\rho$  gives the claimed estimate.  $\square$

### 3.3 Critical inequality for $E_\theta$

**Lemma 3.3.** *For all  $\rho > 1$ ,*

$$E_\theta(\rho) \leq C_\theta \rho^{-1} \left( \|\nabla \mathbf{u}\|_{L^2(A(\theta\rho, \rho))} \|\mathbf{u}\|_{L^2(A(\theta\rho, \rho))} + \|\nabla \mathbf{b}\|_{L^2(A(\theta\rho, \rho))} \|\mathbf{b}\|_{L^2(A(\theta\rho, \rho))} \right) \\ + C_\theta \rho^{-1} \left( \|\mathbf{u}\|_{L^3(A(\theta\rho, \rho))}^3 + \|\mathbf{b}\|_{L^3(A(\theta\rho, \rho))}^3 \right).$$

*Proof.* Fix  $\rho > 1$  and let  $S = A(\theta\rho, \rho)$ . Choose a smooth radial cut-off  $\psi \in C_c^\infty(S)$  such that  $\psi \equiv 1$  on  $A(\theta'\rho, \rho')$  with  $\theta < \theta' < \rho'/\rho < 1$ , and  $|\nabla\psi| \lesssim \rho^{-1}$ ,  $|\nabla^2\psi| \lesssim \rho^{-2}$ .

To eliminate the pressure, we use the Bogovskii operator on  $S$ . Since  $\nabla \cdot \mathbf{u} = 0$ ,  $\int_S \mathbf{u} \cdot \nabla(\psi^2) = 0$ , so  $f := \mathbf{u} \cdot \nabla(\psi^2) \in L^q_0(S)$ . Set  $\mathbf{w} = Tf$ ; then  $\nabla \cdot \mathbf{w} = f$  and  $\|\nabla \mathbf{w}\|_{L^q(S)} \lesssim \rho^{-1} \|\mathbf{u}\|_{L^q(S)}$  for any  $q \in (1, \infty)$ .

Test the momentum equation with  $\mathbf{u}\psi^2 - \mathbf{w}$  and the induction equation with  $\mathbf{b}\psi^2$ . Repeating the integration by parts of Lemma 3.1 on  $S$  (all boundary terms vanish because  $\psi$  and  $\mathbf{w}$  are supported in  $S$ ) we obtain exactly the same algebraic identity as (6) with  $\phi$  replaced by  $\psi$  and  $\mathbf{v}$  by  $\mathbf{w}$ .

Estimating all terms with the bounds for  $\psi$  and  $\mathbf{w}$  and letting  $\theta' \rightarrow \theta$ ,  $\rho' \rightarrow \rho$  yield the stated inequality. The details are analogous to the scalar case treated in [4].  $\square$

### 3.4 Growth condition implies gradient bound

**Lemma 3.4.** *Assume that for some  $\frac{3}{2} < p < 3$  and all large  $\rho$ ,*

$$\|\mathbf{u}\|_{L^p(A(\theta\rho, \theta^{-1}\rho))} + \|\mathbf{b}\|_{L^p(A(\theta\rho, \theta^{-1}\rho))} \leq \rho^{\frac{2}{p}-\frac{1}{3}} g(\rho)^{\frac{3}{p}-1}.$$

*Then*

$$\|\mathbf{u}\|_{L^6(B(\rho))} + \|\mathbf{b}\|_{L^6(B(\rho))} + \|\nabla \mathbf{u}\|_{L^2(B(\rho))} + \|\nabla \mathbf{b}\|_{L^2(B(\rho))} \leq C_\theta g(\rho)^{\frac{9-3p}{4p-6}}.$$

*Proof.* Substitute the hypothesis into Lemma 3.2. Since  $g(\rho) \geq 1$ ,

$$\rho^{-\frac{6-p}{2p-3}} \left( \rho^{\frac{2}{p}-\frac{1}{3}} g(\rho)^{\frac{3}{p}-1} \right)^{\frac{3p}{2p-3}} = g(\rho)^{\frac{9-3p}{2p-3}}.$$

The  $\rho^{-1}$  term is bounded by the same quantity. Taking square roots gives the result.  $\square$

## 4 Proof of Theorem 1.1

By Lemma 3.1(i) it suffices to prove  $\liminf_{\rho \rightarrow \infty} K_\theta(\rho) = 0$ . From the Gagliardo–Nirenberg inequality,

$$K_\theta(\rho)^{1/3} \lesssim \rho^{-1/3} \left( \|\mathbf{u}\|_{L^p}^{\frac{p}{6-p}} \|\nabla \mathbf{u}\|_{L^2}^{\frac{6-2p}{6-p}} + \|\mathbf{b}\|_{L^p}^{\frac{p}{6-p}} \|\nabla \mathbf{b}\|_{L^2}^{\frac{6-2p}{6-p}} \right) \\ + \rho^{\frac{2}{3}-\frac{3}{p}} \left( \|\mathbf{u}\|_{L^p} + \|\mathbf{b}\|_{L^p} \right),$$

with norms over  $A = A(\theta\rho, \theta^{-1}\rho)$ . Hypothesis (2) gives

$$\|\mathbf{u}\|_{L^p} + \|\mathbf{b}\|_{L^p} \lesssim \rho^{\frac{2}{p}-\frac{1}{3}} g(\rho)^{\frac{3}{p}-1}.$$

Thus

$$K_\theta(\rho)^{1/3} \lesssim (g(\rho) \|\nabla \mathbf{u}\|_{L^2}^2)^{\frac{3-p}{6-p}} + (g(\rho) \|\nabla \mathbf{b}\|_{L^2}^2)^{\frac{3-p}{6-p}} + (\rho^{-1/3} g(\rho))^{\frac{3}{p}-1}.$$

The last term tends to 0 by Assumption 1.3(2). Now suppose that  $\liminf_{\rho \rightarrow \infty} K_\theta(\rho) > 0$ . Then there exist  $\delta > 0$  and  $\rho_0 \geq 1$  such that  $K_\theta(\rho) \geq \delta$  for all  $\rho \geq \rho_0$ . Hence, for large  $\rho$ ,

$$\delta^{1/3} \leq C(g(\rho)\|\nabla \mathbf{u}\|_{L^2(A)}^2)^{\frac{3-p}{6-p}} + C(g(\rho)\|\nabla \mathbf{b}\|_{L^2(A)}^2)^{\frac{3-p}{6-p}} + o(1).$$

Consequently, there exists  $c_0 > 0$  such that

$$F(\rho) := \|\nabla \mathbf{u}\|_{L^2(A)}^2 + \|\nabla \mathbf{b}\|_{L^2(A)}^2 \geq \frac{c_0}{g(\rho)}. \quad (9)$$

Now we use the energy  $E_\theta$ . From Lemma 2.2,

$$E'_\theta(\rho) \geq \frac{\theta}{\rho(1-\theta)}F(\rho) \geq \frac{c_1}{\rho g(\rho)}, \quad c_1 > 0.$$

Fix  $\rho_1 \geq \rho_0$  and consider the dyadic radii  $\rho_k = \theta^{-2k}\rho_1$ . Integrating the differential inequality,

$$E_\theta(\rho_k) - E_\theta(\rho_1) \geq c_1 \int_{\rho_1}^{\rho_k} \frac{d\rho}{\rho g(\rho)}.$$

Letting  $k \rightarrow \infty$ , the integral diverges because  $\int^\infty \frac{d\rho}{\rho g(\rho)} = \infty$  by Assumption 1.3(3). Hence  $E_\theta(\rho_k) \rightarrow \infty$ .

However, by Theorem 1.1's hypothesis we have  $\nabla \mathbf{u}, \nabla \mathbf{b} \in L^2(\mathbb{R}^3)$ . Thus

$$E_\theta(\rho) = \int_{\mathbb{R}^3} (|\nabla \mathbf{u}|^2 + |\nabla \mathbf{b}|^2)\eta_\theta(|x|/\rho)dx \leq \|\nabla \mathbf{u}\|_{L^2}^2 + \|\nabla \mathbf{b}\|_{L^2}^2 < \infty,$$

which contradicts  $E_\theta(\rho_k) \rightarrow \infty$ . Therefore  $\liminf K_\theta(\rho) = 0$ , completing the proof.

## 5 Proof of Theorem 1.2

**Lemma 5.1.** *Let  $g$  satisfy Assumption 1.3. For any  $0 < \theta < 1$  and  $3/2 < p < 3$ ,*

$$\liminf_{\rho \rightarrow \infty} \frac{\|\mathbf{u}\|_{L^3(A)} + \|\mathbf{b}\|_{L^3(A)}}{\rho^{1/3}} \leq \limsup_{\rho \rightarrow \infty} \frac{\|\mathbf{u}\|_{L^p(A)} + \|\mathbf{b}\|_{L^p(A)}}{\rho^{\frac{2}{p}-\frac{1}{3}}g(\rho)^{\frac{3}{p}-1}},$$

with  $A = A(\theta\rho, \theta^{-1}\rho)$ .

*Proof.* Assume the contrary, i.e. there exist  $M > 0$  and  $\rho_0 \geq 1$  such that for all  $\rho \geq \rho_0$ ,

$$\|\mathbf{u}\|_{L^p} + \|\mathbf{b}\|_{L^p} < M\rho^{\frac{2}{p}-\frac{1}{3}}g(\rho)^{\frac{3}{p}-1} < (\|\mathbf{u}\|_{L^3} + \|\mathbf{b}\|_{L^3})\rho^{\frac{2}{p}-\frac{2}{3}}g(\rho)^{\frac{3}{p}-1}.$$

From the second inequality we obtain, after canceling the common factor  $\rho^{\frac{2}{p}}g(\rho)^{\frac{3}{p}-1}$ ,

$$\|\mathbf{u}\|_{L^3(A)} + \|\mathbf{b}\|_{L^3(A)} > M\rho^{1/3}. \quad (10)$$

Let  $\sigma = \theta^2$  and  $A_\sigma = A(\sigma\rho, \rho)$ . Lemma 3.3 together with Young's inequality gives

$$\begin{aligned} E_\sigma(\rho) &\leq \epsilon\rho^{-1}(\|\nabla \mathbf{u}\|_{L^2(A_\sigma)}^2 + \|\nabla \mathbf{b}\|_{L^2(A_\sigma)}^2) \\ &\quad + C_{\epsilon,\theta}\rho^{-1}(\|\mathbf{u}\|_{L^2(A_\sigma)}^2 + \|\mathbf{b}\|_{L^2(A_\sigma)}^2) \\ &\quad + C_\theta\rho^{-1}(\|\mathbf{u}\|_{L^3(A_\sigma)}^3 + \|\mathbf{b}\|_{L^3(A_\sigma)}^3). \end{aligned}$$

By Lemma 2.2,  $\|\nabla \mathbf{u}\|_{L^2(A_\sigma)}^2 + \|\nabla \mathbf{b}\|_{L^2(A_\sigma)}^2 \leq \frac{\rho(1-\sigma)}{\sigma} E'_\sigma(\rho)$ . Moreover,  $\|f\|_{L^2(A_\sigma)}^2 \leq |A_\sigma|^{1/3} \|f\|_{L^3(A_\sigma)}^2 \leq C\rho \|f\|_{L^3}^2$  and  $\rho^{-1} \cdot \rho \|f\|_{L^3}^2 = \|f\|_{L^3}^2 \leq \eta \|f\|_{L^3}^3 + C_\eta$ . Choosing  $\epsilon, \eta$  sufficiently small, we can absorb the gradient and  $L^2$  terms into the left-hand side and the cubic term. Hence

$$E_\sigma(\rho) \leq C\rho E'_\sigma(\rho) + C\rho^{-1} (\|\mathbf{u}\|_{L^3(A)}^3 + \|\mathbf{b}\|_{L^3(A)}^3). \quad (11)$$

We now relate the cubic term to  $E'_\sigma$  and  $g$ . By Gagliardo–Nirenberg and Sobolev,

$$\|\mathbf{u}\|_{L^3} \leq C \|\mathbf{u}\|_{L^p}^{\frac{p}{6-p}} \|\mathbf{u}\|_{L^6}^{\frac{6-2p}{6-p}} \leq C \|\mathbf{u}\|_{L^p}^{\frac{p}{6-p}} (\|\nabla \mathbf{u}\|_{L^2} + \rho^{-1} \|\mathbf{u}\|_{L^2})^{\frac{6-2p}{6-p}}.$$

The lower-order term can be absorbed for large  $\rho$ . Using the growth hypothesis on the  $L^p$  norms and Lemma 2.2,

$$\begin{aligned} \rho^{-1} \|\mathbf{u}\|_{L^3}^3 &\leq C\rho^{-1} (\rho^{\frac{2}{p}-\frac{1}{3}} g(\rho)^{\frac{3}{p}-1})^{\frac{3p}{6-p}} (\rho E'_\sigma(\rho))^{\frac{3(3-p)}{6-p}} \\ &\leq Cg(\rho)^{-\frac{3(3-p)}{6-p}} (\rho E'_\sigma(\rho))^{\frac{3(3-p)}{6-p}}. \end{aligned}$$

Set  $\alpha = \frac{3(3-p)}{6-p} \in (0, 1)$ . For any  $\delta > 0$ , Young's inequality yields

$$g(\rho)^{-\alpha} (\rho E'_\sigma)^{\alpha} \leq \delta \rho E'_\sigma + C_\delta g(\rho)^{-\frac{\alpha}{1-\alpha}}.$$

Since  $g(\rho) \rightarrow \infty$ , the last term is bounded by a constant. Choosing  $\delta$  small and inserting the resulting estimate for the cubic term into (11), we obtain

$$E_\sigma(\rho) \leq C_1 \rho E'_\sigma(\rho) + C_2. \quad (12)$$

On the other hand, from (10) we have the lower bound  $\rho^{-1} (\|\mathbf{u}\|_{L^3}^3 + \|\mathbf{b}\|_{L^3}^3) \geq M^3/2$ . Together with (11) this implies  $E_\sigma(\rho) \geq M^3/4$  for large  $\rho$ . Combining with (12) we deduce

$$\rho E'_\sigma(\rho) \geq cE_\sigma(\rho) - C_2 \geq \frac{c}{2} E_\sigma(\rho)$$

provided  $E_\sigma(\rho)$  is large enough (which is guaranteed by  $M^3/4$  if  $M$  is chosen appropriately). Thus, for all sufficiently large  $\rho$ ,

$$E'_\sigma(\rho) \geq \frac{c}{2\rho} E_\sigma(\rho). \quad (13)$$

Integrating (13) from  $\rho_1$  to  $R$  gives

$$\ln \frac{E_\sigma(R)}{E_\sigma(\rho_1)} \geq \frac{c}{2} \ln \frac{R}{\rho_1} \implies E_\sigma(R) \geq E_\sigma(\rho_1) \left(\frac{R}{\rho_1}\right)^{c/2}.$$

Hence  $E_\sigma(R)$  grows at least polynomially.

But we can also bound  $E_\sigma$  from above using Lemma 3.4 together with the  $L^p$  growth condition:

$$E_\sigma(\rho) \leq Cg(\rho)^{\frac{9-3p}{2p-3}}.$$

Since  $g$  satisfies  $\lim_{\rho \rightarrow \infty} \rho^{-1/3} g(\rho) = 0$ , we have  $g(\rho) = o(\rho^{1/3})$ , and therefore  $g(\rho)^{\frac{9-3p}{2p-3}} = o(\rho^{\frac{9-3p}{3(2p-3)}})$ . Because  $\frac{9-3p}{3(2p-3)}$  is a fixed positive number (for  $p \in (3/2, 3)$ ), the polynomial growth  $\rho^{c/2}$  will eventually dominate, contradicting the upper bound. This completes the proof.  $\square$

*Proof of Theorem 1.2.* Because the right-hand side of Lemma 5.1 is finite by (2), we have  $\liminf K_\theta(\rho) < \infty$ . Lemma 3.1(ii) gives  $\nabla \mathbf{u}, \nabla \mathbf{b} \in L^2(\mathbb{R}^3)$ , and Theorem 1.1 forces  $\mathbf{u} = \mathbf{b} = 0$ .  $\square$

**Data Availability Statement** Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

**Conflict of interest** On behalf of all authors, the corresponding author states that there is no conflict of interest.

## References

- [1] G. P. Galdi, *An introduction to the mathematical theory of the Navier–Stokes equations. Steady-state problems*, 2nd ed., Springer, New York, 2011.
- [2] G. Seregin, *Liouville type theorem for stationary Navier–Stokes equations*, *Nonlinearity* **29** (2016), no. 8, 2191–2195.
- [3] G. Seregin and W. Wang, *Sufficient conditions on Liouville type theorems for the 3D steady Navier–Stokes equations*, *Algebra i Analiz* **31** (2019), no. 2, 269–278.
- [4] T.-P. Tsai, *Liouville type theorems for stationary Navier–Stokes equations*, *Partial Differ. Equ. Appl.* **2** (2021), no. 1, Paper No. 10, 20 pp.
- [5] D. Chae and J. Wolf, *On Liouville type theorems for the steady Navier–Stokes equations in  $\mathbb{R}^3$* , *J. Differential Equations* **261** (2016), no. 10, 5541–5560.
- [6] D. Chae and J. Wolf, *On Liouville type theorems for the stationary Navier–Stokes equations*, *Calc. Var. Partial Differential Equations* **58** (2019), no. 3, Paper No. 111, 11 pp.
- [7] Y. Cho, J. Neustupa, and M. Yang, *New Liouville type theorems for the stationary Navier–Stokes, MHD, and Hall-MHD equations*, *Nonlinearity* **37** (2024), no. 3, Paper No. 035007, 22 pp.
- [8] H. J. Choe and M. Yang, *Local kinetic energy and singularities of the incompressible Navier–Stokes equations*, *J. Differential Equations* **264** (2018), no. 2, 1171–1191.
- [9] H. Kozono, Y. Terasawa, and Y. Wakasugi, *A remark on Liouville-type theorems for the stationary Navier–Stokes equations in three space dimensions*, *J. Funct. Anal.* **272** (2017), no. 2, 804–818.
- [10] J. Bang and Z. Yang, *Saint-Venant estimates and Liouville-type theorems for the stationary Navier–Stokes equation in  $\mathbb{R}^3$* , *J. Math. Fluid Mech.* **27** (2025), no. 3, Paper No. 35, 13 pp.
- [11] Y. Cho, J. Choi, and M. Yang, *A Liouville-type theorem for the stationary Navier–Stokes equations*, *Appl. Math. Lett.* **143** (2023), Paper No. 108664, 6 pp.
- [12] D. Chae, *Liouville-type theorems for the forced Euler equations and the Navier–Stokes equations*, *Comm. Math. Phys.* **326** (2014), no. 1, 37–48.