

Synergistic Effects on Silicon and Selenium on Growth, Yield and Protein Content in Blackgram (*Vigna mungo* L.) Grown in Coastal Saline Soil

ABSTRACT

Salinity stress severely limits pulse productivity in coastal agro-ecosystems by inducing ionic imbalance and oxidative damage. Although silicon and selenium are known to enhance plant stress tolerance individually, information on their combined application under coastal saline soil conditions is limited. A pot experiment was conducted to evaluate the interactive effects of silicon and selenium on growth, yield and protein content of blackgram (*Vigna mungo* L.). Treatments consisted of graded levels of silicon and selenium applied individually and in combination along with recommended dose of fertilizers. The combined application of selenium at the rate of 2 ppm and silicon at the rate of 50 kg ha⁻¹ significantly improved plant height, dry matter production, nodulation, yield attributes and grain yield (1224 kg ha⁻¹) compared to control. Protein content and protein yield were also maximized under the same treatment. The improved performance may be attributed to enhanced nutrient uptake, improved physiological efficiency and alleviation of salinity-induced stress. The study highlights the synergistic role of silicon and selenium in improving productivity and grain quality of blackgram under coastal saline soil conditions.

Keywords: Blackgram; Silicon; Selenium; Salinity; Coastal saline soil; Abiotic stress

1. INTRODUCTION

Pulses hold significant value in Indian agriculture due to their high protein content, as well as their abundance in vitamins, minerals, and specific fibers. They are famously referred to as the "Poor man's meat" and "rich man's vegetable" [1]. Blackgram (*Vigna mungo* L.) is a pivotal pulse crop in India and South Asia. It holds a key place in Indian agriculture, boasting a high protein content of about 26%, nearly three times that of cereals. Silicon (Si) enhances plant resistance to various stresses like drought, salinity and pests by strengthening cell walls, improving nutrient uptake, and boosting overall plant health and yield [2]. Selenium [Se], a vital micronutrient, promotes plant growth and stress tolerance, improves antioxidant defenses, and mitigates the negative effects of soil pH on plant growth. Absorbed from the soil, selenium can biofortify crops, enhancing their nutritional value for human and animal consumption [3].

Salinity stress, due to high soil salt concentrations, negatively impacts crop growth and productivity, leading to osmotic stress, ion toxicity, and oxidative stress, which result in reduced water uptake, nutrient imbalances, and impaired photosynthesis. Coastal salinity is a major factor leading to poor crop yields over approximately 3.1 million hectares [4]. Coastal saline soils present particular challenges, including a light texture, poor exchange properties,

low nutrient and water retention capacity, low organic carbon levels, and deficiencies in both macro and micronutrients [5].

Recent studies indicate that silicon (Si) and selenium (Se) enhance plant tolerance to salinity stress by improving growth and overall performance. Their combined application has shown synergistic effects in improving plant productivity under stress conditions. However, most studies are limited to cereals and horticultural crops, with very limited information available on pulse crops, particularly blackgram under coastal saline soils. Although the individual effects of Si and Se are well documented, their combined influence on growth, yield attributes, and grain protein content in blackgram under saline conditions remains insufficiently explored [6]. Therefore, the present study was undertaken to evaluate the synergistic effect of silicon and selenium on growth, yield and grain protein content of blackgram under coastal saline soil conditions.

2. MATERIAL AND METHODS

A pot culture experiment was conducted at the pot-culture yard, Department of Soil Science and Agricultural Chemistry, Faculty of Agriculture, Annamalai University, Tamil Nadu, India, during January to April 2024. The experiment was designed to maintain uniform salinity stress conditions and ensure precise nutrient application. The experimental soil was collected from a coastal saline field in Pichavaram village. The soil was sandy loam in texture, low in available nitrogen and phosphorus and medium in available potassium. The experiment consisted of ten treatments laid out in a completely randomized design (CRD) with three replications. The recommended dose of fertilizers (RDF) was applied as per crop recommendation (25:50:25 N:P₂O₅:K₂O kg ha⁻¹). Selenium was applied as sodium selenate (Na₂SeO₄) at concentrations of 1, 2, and 3 ppm, while silicon was applied through diatomaceous earth at rates equivalent to 25, 50, and 75 kg ha⁻¹.

Growth parameters including plant height, dry matter production, leaf area index, and number of branches were recorded at different growth stages. Yield attributes such as number of pods per plant, pod length, number of seeds per pod, 100-seed weight, grain yield, and haulm yield were recorded to assess crop productivity. Seed protein content was calculated by multiplying nitrogen percentage in grain with the conversion factor 6.25 [7]. Soil pH and electrical conductivity (EC) were analyzed to confirm the saline nature of the soil [8].

3. RESULTS AND DISCUSSION

3.1 Growth parameters

The application of silicon and selenium significantly improved the growth parameters of blackgram under coastal saline soil conditions (Table 1). In general, the combined application of silicon and selenium produced a greater response than their individual application, clearly indicating a synergistic effect.

Among the treatments, T9 (RDF + selenium @ 2 ppm + silicon @ 50 kg ha⁻¹) recorded the highest plant height (51.8 cm), number of branches plant⁻¹ (11.25), number of leaves plant⁻¹ (21.6), leaf area index (2.74), chlorophyll content (46.75 SPAD), and number of nodules plant⁻¹ (29.08). This treatment was closely followed by T8 (RDF + selenium @ 1 ppm + silicon @ 25 kg ha⁻¹), which also showed marked improvement over the control and individual nutrient applications. The control treatment (T1) recorded the lowest values for all growth parameters. The enhanced vegetative growth under combined application of silicon and selenium may be attributed to better regulation of ionic balance, improved nutrient uptake, stronger photosynthetic activity and protection against salinity-induced oxidative stress [9]. Silicon likely reduced sodium toxicity and improved tissue structural stability, whereas selenium may have enhanced antioxidant defence and chlorophyll preservation. These combined effects would have favoured higher leaf development, greater nodulation, and increased biomass accumulation. Similar findings have been reported [10,11].

Table 1. Effect of silicon and selenium on growth parameters of blackgram at harvest

Treatments	Plant height (cm)	Number of branches plant ⁻¹	Number of leaves plant ⁻¹	Leaf area index	Chlorophyll content (SPAD)	Number of nodules plant ⁻¹
T1 – RDF (control)	23.4	6.15	11.8	1.01	22.28	16.51
T2 – RDF + Se @ 1 ppm	34.1	7.92	15.3	1.64	30.82	21.25
T3 – RDF + Se @ 2 ppm	30.8	7.35	14.2	1.43	27.96	19.11
T4 – RDF + Se @ 3 ppm	27.5	6.79	13.1	1.22	25.13	17.85
T5 –RDF + Si @ 25 kg ha ⁻¹	37.3	8.49	16.5	1.85	33.65	23.37
T6 – RDF + Si @ 50 kg ha ⁻¹	40.5	9.05	17.6	2.06	36.45	25.48
T7 – RDF + Si @ 75 kg ha ⁻¹	45.4	10.09	19.4	2.31	41.15	26.65
T8 – RDF + Se @ 1 ppm + Si @ 25 kg ha ⁻¹	48.6	10.66	20.5	2.53	43.95	27.88
T9 – RDF + Se @ 2 ppm + Si @ 50 kg ha ⁻¹	51.8	11.25	21.6	2.74	46.75	29.08
T10 – RDF + Se @ 3 ppm + Si @ 75 kg ha ⁻¹	42.2	9.51	18.2	2.10	38.26	25.50
S. Ed	1.39	0.26	0.52	0.09	1.235	0.45
CD (p = 0.05)	3.0	0.55	1.14	0.21	2.71	0.95

3.2 Dry matter production and yield attributes

Dry matter production, yield attributes, and yield of blackgram were significantly influenced by the different treatments (Table 2). The highest dry matter production at harvest was recorded in T9 (35.72 g pot⁻¹), followed by T8 (33.88 g pot⁻¹), whereas the lowest was observed in the control (23.19 g pot⁻¹). Yield-contributing traits followed a similar trend, with T9 recording the highest pod length (8.24 cm), number of pods plant⁻¹ (19.55) and number of seeds pod⁻¹ (7.89). Although 100-seed weight did not differ significantly among treatments, a slight numerical increase was observed under combined application, with T9 recording the highest value (3.58 g).

Grain and haulm yields were also significantly enhanced by silicon and selenium application, with combined treatments outperforming individual applications. T9 recorded the highest grain yield (1224 kg ha⁻¹) and haulm yield (1947 kg ha⁻¹), followed by T8 (1142 and 1876 kg ha⁻¹, respectively), while the control recorded the lowest values (763 and 1290 kg ha⁻¹). Compared with the control, T9 increased grain yield by 60.4% and haulm yield by 50.9%. The superior performance under T9 indicates that the combined application of silicon and selenium improved assimilate production and partitioning, resulting in enhanced vegetative growth and reproductive efficiency under saline conditions. Silicon-alone treatments also performed better than selenium-alone treatments, but the maximum response was observed under combined application, confirming their synergistic effect. These improvements may be attributed to enhanced physiological efficiency, better chlorophyll retention, and increased tolerance to salinity stress, which is in agreement with earlier findings [12].

Table 2. Effect of silicon and selenium on dry matter production, yield attributes and yield of blackgram

Treatments	Dry matter production (g pot ⁻¹)	Pod length (cm)	Pods plant ⁻¹	Seeds pod ⁻¹	100-seed weight (g)	Grain yield (kg ha ⁻¹)	Haulm yield (kg ha ⁻¹)
T1	23.19	4.85	8.04	4.01	3.39	763	1290
T2	27.23	6.02	12.13	5.31	3.46	952	1656
T3	26.88	5.65	10.91	4.89	3.45	935	1591
T4	25.01	5.31	9.55	4.46	3.41	881	1420
T5	28.03	6.41	13.46	5.74	3.48	976	1677
T6	29.91	6.78	14.98	6.18	3.49	1030	1746
T7	31.96	7.41	17.12	6.90	3.53	1078	1806
T8	33.88	7.78	18.23	7.40	3.54	1142	1876
T9	35.72	8.24	19.55	7.89	3.58	1224	1947
T10	30.66	7.05	15.95	6.45	3.53	1052	1788
S. Ed	0.792	0.147	0.504	0.195	0.04	23.49	29.64
CD (p = 0.05)	1.75	0.32	1.10	0.41	NS	53.0	68.0

3.3 Protein content and protein yield

Protein content and protein yield were significantly influenced by the application of silicon and selenium (Table 3). The highest protein content (29.84%) and protein yield (365 kg ha⁻¹) were recorded under T9, followed by T8, which recorded 27.97% protein content and 319 kg ha⁻¹ protein yield. The lowest values were recorded in the control treatment, with 15.94% protein content and 121 kg ha⁻¹ protein yield.

The increase in protein content under combined application may be attributed to improved nitrogen assimilation and enhanced metabolic activity in plants. Selenium is known to promote protein synthesis through its role in nitrogen metabolism and antioxidant protection, whereas silicon improves nutrient absorption and reduces salinity-induced physiological injury [13,14]. The combined application of these nutrients therefore created a favourable environment for greater protein accumulation in grain. The increase in protein yield under T9 was further supported by the substantial rise in grain yield. Similar improvements in grain quality with selenium and silicon application have been reported by [15,16].

Table 3. Effect of silicon and selenium on protein content and protein yield of blackgram

Treatments	Protein content (%)	Protein yield (kg ha ⁻¹)
T1	15.94	121
T2	21.40	203
T3	20.78	194
T4	18.90	166
T5	22.03	215
T6	23.75	244
T7	26.09	281
T8	27.97	319
T9	29.84	365
T10	25.00	263
S. Ed	0.763	8.085
CD (p = 0.05)	1.60	17.0

4. CONCLUSION

The present study demonstrated that the integrated application of silicon and selenium significantly improved the growth, yield, and protein content of blackgram grown in coastal saline soil. Among the treatments, RDF + selenium @ 2 ppm + silicon @ 50 kg ha⁻¹ (T9) proved to be the most effective, recording the highest values for plant growth, nodulation, chlorophyll content, dry matter production, yield attributes, grain yield, haulm yield, protein content, and protein yield.

The results confirm that silicon and selenium act synergistically in mitigating salinity stress and enhancing both productivity and grain quality of blackgram. Therefore, the integrated use of selenium @ 2 ppm and silicon @ 50 kg ha⁻¹ along with the recommended dose of fertilizers may be considered a promising nutrient management strategy for blackgram cultivation in coastal saline soils. However, as the present investigation was conducted under pot culture conditions, further validation through multi-location field trials in different coastal saline agro-ecological regions is necessary before making broad-scale recommendations.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that they have no known competing financial interests OR non-financial interests OR personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

1. Singh, A. K., Singh, S. S., Prakash, V. E. D., Kumar, S. and Dwivedi, S. K. Pulses production in india: Present status, sent status, bottleneck and way forward. Journal of AgriSearch. (2015);2(2):p.75-83.
2. Malik, M. A., Wani, A. H., Mir, S. H., Rehman, I. U., Tahir, I., Ahmad, P and Rashid, I. Elucidating the role of silicon in drought stress tolerance in plants. Plant Physiology and Biochemistry. (2021);165:p.187-195.
3. Jawad Hassan, M., Ali Raza, M., Khan, I., Ahmad Meraj, T., Ahmed, M., Abbas Shah, G and Li, Z. Selenium and salt interactions in black gram (*Vigna mungo* L.): ion uptake, antioxidant defense system, and photochemistry efficiency. Plants. (2020);9(4):p.467.
4. Arulmathi, C. and Porkodi, G. Characteristics of coastal saline soil and their management: A review. International Journal of Current Microbiology and Applied Sciences. (2020); 9(10):p.1726-1734.
5. Sekaran, D., Singaravel, R and Valavan, P. S. Effect of micronutrient and bioactive compounds fortified organic manure on the growth and yield of blackgram in coastal saline soil. Research on Crops.(2020);21(1):p.54-58.

6. Xu, S., Zhao, N., Qin, D., Liu, S., Jiang, S., Xu, L., ... & Hu, A. (2021). The synergistic effects of silicon and selenium on enhancing salt tolerance of maize plants. *Environmental and Experimental Botany*, 187, 104482.
7. Piper, C.S. Soil and plant analysis, Hans Publishers, Bombay, India; 1966.
8. Jackson, M. L. Soil chemical analysis. Prentice Hall of India Pvt. Ltd., New Delhi;1973.
9. Taha, R. S., Seleiman, M. F., Shami, A., Alhammad, B. A., & Mahdi, A. H. (2021). Integrated application of selenium and silicon enhances growth and anatomical structure, antioxidant defense system and yield of wheat grown in salt-stressed soil. *Plants*, 10(6), 1040.
10. Berahim, Z., Omar, M. H., Zakaria, N. I., Ismail, M. R., Rosle, R., Roslin, N. A., and Cheya, N. N. Silicon improves yield performance by enhancement in physiological responses, crop imagery, and leaf and culm sheath morphology in new rice line, Padiu Putra. *BioMed research international*. (2021);1:p.6679787.
11. Hossain, A., Skalicky, M., Brestic, M., Maitra, S., Sarkar, S., Ahmad, Z., and Laing, A. M. Selenium biofortification: roles, mechanisms, responses and prospects. *Molecules*. (2021);26(4):p.881.
12. Hu, W., Su, Y., Zhou, J., Zhu, H., Guo, J., Huo, H., and Gong, H. Foliar application of silicon and selenium improves the growth, yield and quality characteristics of cucumber in field conditions. *Scientia Horticulturae*. (2022);294:p.110776.
13. Khan, Z., Thounaojam, T. C., Chowdhury, D., & Upadhyaya, H. (2023). The role of selenium and nano selenium on physiological responses in plant: a review. *Plant Growth Regulation*, 100(2), 409-433.
14. Al Murad, M., & Muneer, S. (2022). Silicon supplementation modulates physiochemical characteristics to balance and ameliorate salinity stress in mung bean. *Frontiers in Plant Science*, 13, 810991.
15. Nawaz, F., Ashraf, M. Y., Ahmad, R., Waraich, E. A., Shabbir, R. N., and Bukhari, M. A. Supplemental selenium improves wheat grain yield and quality through alterations in biochemical processes under normal and water deficit conditions. *Food Chemistry*. (2015); 175:p.350-357.
16. Lara, T. S., Correia, T. S., de Oliveira, C., Lessa, J. H. D. L., de Souza, K. R. D., Corguinha, A. P. B., ... & Guilherme, L. R. G. (2024). Selenium application provides nutritional and metabolic benefits to wheat plants. *Agronomy*, 14(3), 462.